

Exact ground state for 1D electronic models

D.V.Dmitriev ¹, V.Ya.Krivnov ² and A.A.Ovchinnikov ^{1,2}

¹ Max-Planck-Institut für Physik Komplexer Systeme,
Nothnitzer Str. 38, 01187 Dresden, Germany

² Joint Institute of Chemical Physics of RAS,
Kosygin str.4, 117977, Moscow, Russia

Abstract

We have found the exact ground state for two electronic models on a linear chain. The first model describes half-filling electron system in the ferromagnet–antiferromagnet transition point. In the singlet ground state the spin correlators show giant spiral magnetic ordering with the period of a spiral equals to the system size. The second electronic model describes the point where the ground state has giant spiral off-diagonal long-range order and, therefore, superconducting. We suggest the formation of the ground state with giant spiral order (ferromagnetic or off-diagonal) as a probable scenario of the subsequent destruction of the ferromagnetism and the superconductivity.

1 Introduction

In recent years there has been a growing interest in studying systems of strongly correlated electrons in relation with high- T_c superconductivity. Because of the difficulty in dealing with the many-body problems the exact results are rare. It is well-known that some 1D electron models can be exactly solved by the Bethe ansatz. However, a lot of 1D quantum systems do not obey the Yang-Baxter equation, and thus are non-integrable. Another approach leading to exact results consists in construction of the exact ground state wave function for some quantum systems. Recently considerable progress in this problem has been achieved by using the so-called matrix-product (MP) form of the ground state wave function. It allowed to find exact ground state for various 1D spin models [1]-[3]. Its origin can be traced back to the $S = 1$ spin chain model [4]. For higher dimensional spin and electronic systems, there are also some methods for the construction of the exact ground state wave function [5]-[8].

There is a class of the 1D quantum spin models describing the ferromagnet – antiferromagnet transition point, for which the exact ground state wave function has been found in [9, 10]. The singlet ground state wave function at this point has a special recurrent form, and for special values of model parameters it can be reduced to the MP form or to the RVB form [9]. Spin correlations in the singlet ground state show giant spiral magnetic structure with the period of the spiral equals to the system size. On the antiferromagnet side of this point the ground state can be either gapless with the algebraic decay of spin

correlations [11] or gapped with the exponential decay of correlations [12]. So, this model describes the boundary between the ferromagnetic phase and the singlet phase without long-range order.

In this paper we present the singlet ground state wave function of this spin model in another form, which can be easily generalized for the 1D electronic models. Then we consider two 1D electronic models. The first model describes half-filling electron system at the point where the singlet and ferromagnetic states are degenerate. The exact calculation of the correlation functions in the singlet ground state shows the same giant spiral magnetic ordering as for the original spin model, while all other correlations vanish in the thermodynamic limit. The second electronic model describes the boundary on the phase diagram between the superconducting phase with off-diagonal long-range order (ODLRO) and the non-superconducting phase. The correlation functions in the ground state of this model show the giant spiral off-diagonal long-range order. We presume that in one-dimensional systems the destruction of the long-range order (ferromagnetic or off-diagonal) can be followed by the appearance of the ground state with the giant spiral order.

We generalize this form of the wave function for the electronic ladder model. This model possesses both the giant spiral spin order and the giant spiral ODLRO in the ground state. Therefore, this electron ladder model describes the boundary points on the phase diagram between four different phases: two singlet phases with and without ODLRO, and two ferromagnetic phases with and without ODLRO. For some special cases the ground state wave function can be reduced to the usual MP form.

The paper is organized as follows. In Section 2 we construct the exact singlet ground state for the quantum spin model. In Section 3 two electronic models with exact ground states are considered, and the correlation functions are exactly calculated. Section 4 gives a brief summary. In Appendix a technique for the calculation of correlators is developed.

2 Quantum spin model

First, we consider the $s = \frac{1}{2}$ spin chain model with nearest- and next-nearest neighbor interactions given by the Hamiltonian

$$H = - \sum_{i=1}^N (\mathbf{S}_i \cdot \mathbf{S}_{i+1} - \frac{1}{4}) + \frac{1}{4} \sum_{i=1}^N (\mathbf{S}_i \cdot \mathbf{S}_{i+2} - \frac{1}{4}) \quad (1)$$

with periodic boundary conditions and even N .

This model describes the ferromagnet–antiferromagnet transition point where the ferromagnetic and singlet states are degenerate. The Hamiltonian (1) has been considered in [9, 10], where the singlet ground state wave function was constructed in two different forms. In this paper we represent another form of this singlet function, which allows us to generalize this function for the electronic model and to develop a technique to calculate correlators.

The singlet ground state wave function for the Hamiltonian (1) can be written as follows:

$$\Psi_0 = P_0 \Psi, \quad \Psi = \langle 0_b | g_1 \otimes g_2 \otimes \dots \otimes g_N | 0_b \rangle \quad (2)$$

where

$$g_i = b^+ |\uparrow\rangle_i + b |\downarrow\rangle_i \quad (3)$$

Here we introduced one auxiliary Bose-particle b^+ , b (the Bose operators b^+ , b do not act on spin states $|\uparrow\rangle_i$ and $|\downarrow\rangle_i$) and the Bose vacuum $|0_b\rangle$. Therefore, the direct product $g_1 \otimes g_2 \otimes \dots \otimes g_N$ is the superposition of all possible spin configurations multiplied on the corresponding Bose operators, like $b^+ b b b^+ \dots |\uparrow\downarrow\uparrow\dots\rangle$. P_0 is a projector onto the singlet state. This operator can be written as [13]

$$P_0 = \frac{1}{8\pi^2} \int_0^{2\pi} d\alpha \int_0^{2\pi} d\beta \int_0^\pi \sin \gamma d\gamma e^{i\alpha S^z} e^{i\gamma S^x} e^{i\beta S^z}, \quad (4)$$

where $S^{x(z)}$ are components of the total spin operator.

This form of wave function resembles the MP form but with infinity matrix, which is represented by Bose operators. Therefore, we have to pick out the $\langle 0_b | \dots | 0_b \rangle$ element of the matrix product instead of usual *Trace* in the MP formalism [1]-[3], because *Trace* is undefined in this case. The function Ψ contains components with all possible values of spin S ($0 \leq S \leq N/2$) and, in fact, a fraction of the singlet is exponentially small at large N . This component is filtered out by the operator P_0 .

In order to show that Ψ_0 is the ground state wave function for the Hamiltonian (1), let us represent the Hamiltonian (1) as a sum of Hamiltonians h_i of cells containing three sites

$$H = \sum_{i=1}^N h_i, \quad (5)$$

where

$$h_i = -\frac{1}{2}(\mathbf{S}_i \mathbf{S}_{i+1} - \frac{1}{4}) - \frac{1}{2}(\mathbf{S}_{i+1} \mathbf{S}_{i+2} - \frac{1}{4}) + \frac{1}{4}(\mathbf{S}_i \mathbf{S}_{i+2} - \frac{1}{4})$$

One can easily check that each cell Hamiltonian h_i for $i = 1, \dots, N-2$ gives zero when acting on $g_i \otimes g_{i+1} \otimes g_{i+2}$.

Since each h_i is non-negatively defined operator, then Ψ is the exact ground state wave function of an open chain:

$$H_{\text{open}} = \sum_{i=1}^{N-2} h_i$$

As it was mentioned above, the function Ψ contains components with all possible values of total spin S , and, therefore, the ground state of the open chain is multiply degenerate. But it can be proven (as it was done in [10]), that for the cyclic chain (1) only singlet and ferromagnetic components of Ψ have zero energy. Therefore, for the cyclic chain (1) Ψ_0 is the singlet ground state wave function degenerate with the ferromagnetic state.

The exact calculation of the norm and spin correlation function $\langle \mathbf{S}_i \mathbf{S}_{i+l} \rangle$ (see Appendix) in the singlet ground state (2) results in the following expressions:

$$\langle \Psi_0 | \Psi_0 \rangle = \frac{d^N}{d\xi^N} \left(\frac{1}{\cos^2(\frac{\xi}{2})} \right) \Big|_{\xi=0} \quad (6)$$

$$\langle \Psi_0 | \mathbf{S}_i \mathbf{S}_{i+l} | \Psi_0 \rangle = \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \left(-\frac{3 \cos(\xi - \zeta)}{8 \cos^4(\frac{\xi + \zeta}{2})} \right) \Big|_{\xi=\zeta=0} \quad (7)$$

It can be shown that in the thermodynamic limit equations (6) and (7) result in

$$\langle \mathbf{S}_i \mathbf{S}_{i+l} \rangle = \frac{1}{4} \cos \left(\frac{2\pi l}{N} \right) \quad (8)$$

So, we reproduce the result obtained in [9, 10] that in the thermodynamic limit a giant spiral spin structure is realized, with the period of spiral equal to the system size.

3 Electronic models

Now we will construct electronic models by generalization of the wave function (2):

$$\Psi_0 = P_{S=0} \text{Tr}_f \langle 0_b | g_1 \otimes g_2 \otimes \dots \otimes g_N | 0_b \rangle \quad (9)$$

where

$$g_i = b^+ |\uparrow\rangle_i + b |\downarrow\rangle_i + c (f^+ |2\rangle_i + f |0\rangle_i) \quad (10)$$

with b^+ , b and f^+ , f are the Bose and the Fermi operators respectively, $|0_b\rangle$ is the Bose vacuum, and c is a parameter of the model. Here we also denote an empty site by $|0\rangle$, a site occupied by one electron by $|\uparrow\rangle$ and $|\downarrow\rangle$ and a doubly occupied site by $|2\rangle$. So, the product $g_1 \otimes \dots \otimes g_N$ is the operator in the Bose b^+ , b and the Fermi f^+ , f spaces. We pick out the $\langle 0_b | \dots | 0_b \rangle$ element in the Bose space, which can be written as:

$$\langle 0_b | g_1 \otimes g_2 \otimes \dots \otimes g_N | 0_b \rangle = \phi_0 + (f^+ f - f f^+) \phi_1 \quad (11)$$

and then we take the *Trace* over the Fermi operators:

$$\text{Tr}_f \langle 0_b | g_1 \otimes g_2 \otimes \dots \otimes g_N | 0_b \rangle = 2\phi_0$$

The projector $P_{S=0}$ filters out the singlet component from the function ϕ_0 .

Thus, we have a singlet wave function Ψ_0 describing the state with one electron per site.

In order to find the Hamiltonian for which the wave function (9) is the exact ground state wave function, let us consider what states are present on the two nearest sites in the wave function (9). One can easily check that there are only 9 states from the total 16 states in the product $g_i \otimes g_{i+1}$. They are

$$\begin{aligned} & |\uparrow\uparrow\rangle, \quad |\downarrow\downarrow\rangle, \quad |\uparrow\downarrow + \downarrow\uparrow\rangle, \quad |20 - 02\rangle, \quad |\uparrow\downarrow - \downarrow\uparrow\rangle - c^2 |20 + 02\rangle, \\ & |\uparrow 0 - 0 \uparrow\rangle, \quad |\uparrow 2 - 2 \uparrow\rangle, \quad |\downarrow 0 - 0 \downarrow\rangle, \quad |\downarrow 2 - 2 \downarrow\rangle \end{aligned} \quad (12)$$

The elementary Hamiltonian $h_{i,i+1}$ for which all these states are the exact ground states can be written as the sum of the projectors onto the 7 missing states $|\varphi_k\rangle$ with arbitrary positive coefficients λ_k

$$h_{i,i+1} = \sum_{k=1}^7 \lambda_k |\varphi_k\rangle \langle \varphi_k|$$

For $c > 1$ the most simple form of this Hamiltonian is:

$$\begin{aligned} H &= \sum_{i=1}^N h_{i,i+1} \\ h_{1,2} &= 1 - 4 \mathbf{S}_1 \cdot \mathbf{S}_2 + 4 \left(1 - \frac{3}{c^4}\right) \eta_1^z \eta_2^z + \frac{4}{c^4} \eta_1 \cdot \eta_2 \\ &+ \sum_{\sigma} (c_{1,\sigma}^+ c_{2,\sigma} + c_{2,\sigma}^+ c_{1,\sigma}) (1 - n_{1,-\sigma} - n_{2,-\sigma}) \\ &+ \frac{2}{c^2} \sum_{\sigma} (c_{1,\sigma}^+ c_{2,\sigma} + c_{2,\sigma}^+ c_{1,\sigma}) (n_{1,-\sigma} - n_{2,-\sigma})^2 \end{aligned} \quad (13)$$

Here $c_{i,\sigma}^+$, $c_{i,\sigma}$ are the Fermi operators, $n_{i,\sigma} = c_{i,\sigma}^+ c_{i,\sigma}$, and the $SU(2)$ spin operators are given by $S_i^+ = c_{i,\uparrow}^+ c_{i,\downarrow}$, $S_i^- = c_{i,\downarrow}^+ c_{i,\uparrow}$ and $S_i^z = \frac{1}{2}(n_{i,\uparrow} - n_{i,\downarrow})$. We also use here η operators:

$$\eta_i^+ = c_{i,\downarrow}^+ c_{i,\uparrow}^+, \quad \eta_i^- = c_{i,\uparrow} c_{i,\downarrow}, \quad \eta_i^z = \frac{1 - n_{i,\uparrow} - n_{i,\downarrow}}{2},$$

which form another $SU(2)$ algebra [15, 16], and $\eta_1 \cdot \eta_2$ is a scalar product of pseudo-spins η_1 and η_2 .

The Hamiltonian (13) does not conserve the total number of empty and doubly occupied sites because of the last term in the elementary Hamiltonians $h_{i,i+1}$, in contrast to the models considered in [16].

Each elementary Hamiltonian $h_{i,i+1}$ ($i = 1, \dots, N-1$) acting on functions ϕ_0 , ϕ_1 gives zero, since all the states (12) are the eigenstates of $h_{i,i+1}$ with zero energy, while the energies of all other states at $c > 1$ are positive. Therefore, the functions ϕ_0 and ϕ_1 are the ground state wave functions of the open chain:

$$H_{\text{open}} = \sum_{i=1}^{N-1} h_{i,i+1} \quad (14)$$

To determine the degeneracy of the model (14), we need to classify the functions ϕ_n . Analogously to the spin model (2), the functions ϕ_0 and ϕ_1 contain components with all possible values of total spin S . Therefore, ϕ_0 contains multiplets with $S = 0, \dots, N/2$ and ϕ_1 contains components with values of the total spin $S = 0, \dots, N/2 - 1$ (ϕ_1 does not contain the ferromagnetic component, since at least two sites in ϕ_1 are non-magnetic $|0\rangle$ and $|2\rangle$). So, $N + 1$ multiplets are degenerated for the open chain.

But for the cyclic model (13) it can be proved that only three multiplets are the ground states: singlet state (9) with the momentum $p = 0$ (singlet component of ϕ_0), the trivial ferromagnetic state with $p = \pi$, and the state with $S = N/2 - 1$ and $p = \pi$ (which is the component of ϕ_1 with $S = N/2 - 1$). The last state with $S = N/2 - 1$ can be written as:

$$\Psi_{N/2-1} = \sum_{i < j} (c_{i,\uparrow}^+ c_{j,\downarrow} + c_{j,\uparrow}^+ c_{i,\downarrow}) \prod_{n=1}^N c_{n,\downarrow}^+ |0\rangle$$

Thus, the ground states of the electronic model (13) with one electron per site are the singlet state, the ferromagnetic state, and the state with $S = S_{\text{max}} - 1$.

It is interesting to note that the singlet wave function (9) can be also written in the form:

$$\Psi_0 = \sum [i, j][k, l][m, n] \dots \prod_{n=1}^N c_{n,\downarrow}^+ |0\rangle$$

where

$$[i, j] = S_i^+ - S_j^+ + c^2 (c_{i,\uparrow}^+ c_{j,\downarrow} - c_{j,\uparrow}^+ c_{i,\downarrow}),$$

and the summation is made for any combination of sites under the condition that $i < j$, $k < l$, $m < n \dots$. This form of the wave function is analogous to the RVB form found in [9] for the spin model (1).

The norm and the correlators of the electronic model (13) in the singlet ground state are calculated in the same way as for the spin model (Appendix). Therefore, here we give

the final results:

$$\langle \Psi_0 | \Psi_0 \rangle = \frac{d^N}{d\xi^N} \left(2 \frac{1 + \cosh(c^2 \xi)}{\cos^2(\frac{\xi}{2})} \right) \Big|_{\xi=0} \quad (15)$$

$$\langle \Psi_0 | \mathbf{S}_i \mathbf{S}_{i+l} | \Psi_0 \rangle = \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \left(-\frac{3 \cos(\xi - \zeta) (1 + \cosh(c^2 \xi + c^2 \zeta))}{4 \cos^4(\frac{\xi + \zeta}{2})} \right) \Big|_{\xi=\zeta=0} \quad (16)$$

$$\langle \Psi_0 | c_{i,\sigma}^+ c_{i+l,\sigma} | \Psi_0 \rangle = \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \left(-c^2 \frac{(\cos(\xi) + \cos(\zeta)) (\cosh(c^2 \xi) + \cosh(c^2 \zeta))}{2 \cos^4(\frac{\xi + \zeta}{2})} \right) \Big|_{\xi=\zeta=0} \quad (17)$$

$$\langle \Psi_0 | \eta_i^z \eta_{i+l}^z | \Psi_0 \rangle = \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \left(-\frac{c^4 \cosh(c^2 \xi - c^2 \zeta)}{2 \cos^2(\frac{\xi + \zeta}{2})} \right) \Big|_{\xi=\zeta=0} \quad (18)$$

$$\langle \Psi_0 | \eta_i^- \eta_{i+l}^+ | \Psi_0 \rangle = \frac{d^{N-2}}{d\xi^{N-2}} \left(\frac{c^4}{\cos^2(\frac{\xi}{2})} \right) \Big|_{\xi=0} \quad (19)$$

As it can be seen from Eq.(19), the expectation value $\langle \eta_1^- \eta_{l+1}^+ \rangle$, which determines the off-diagonal long-range order (ODLRO) [14], does not depend on the distance l . But in the thermodynamic limit ODLRO vanishes:

$$\begin{aligned} \langle \eta_i^z \eta_{i+l}^z \rangle &= O\left(\frac{1}{N^2}\right), & \langle \eta_i^+ \eta_{i+l}^- \rangle &= O\left(\frac{1}{N^2}\right), \\ \langle c_{i,\sigma}^+ c_{i+l,\sigma} \rangle &= O\left(\frac{1}{N}\right), & \langle \mathbf{S}_i \mathbf{S}_{i+l} \rangle &= \frac{1}{4} \cos\left(\frac{2\pi l}{N}\right), \end{aligned} \quad (20)$$

though for finite systems all correlators (17)-(19) are non zero.

The second electronic model can be obtained by simply interchanging of the Bose and the Fermi operators in (10). So, the wave function of this model has the form:

$$\Psi_0 = P_{\eta=0} Tr_f \langle 0_b | g_1 \otimes g_2 \otimes \dots \otimes g_N | 0_b \rangle \quad (21)$$

with

$$g_i = c (f^+ |\uparrow\rangle_i + f |\downarrow\rangle_i) + b^+ |2\rangle_i + b |0\rangle_i \quad (22)$$

The projector $P_{\eta=0}$ filters out the state with total $\eta = \sum \eta_i = 0$. Therefore, the function Ψ_0 has $S^z = 0$, but it is not an eigenfunction of S^2 . Instead, it is an eigenfunction of η^2 with $\eta = 0$.

The wave function (21) can be also written in the form analogous to the RVB one:

$$\Psi_0 = \sum [i, j][k, l][m, n] \dots |0\rangle$$

where

$$[i, j] = \eta_i^+ - \eta_j^+ + c^2 (c_{i,\uparrow}^+ c_{j,\downarrow}^+ + c_{i,\downarrow}^+ c_{j,\uparrow}^+),$$

and the summation is also done over any combinations of sites under the condition that $i < j, k < l, m < n \dots$

Considering the product $g_i \otimes g_{i+1}$ one can find that there are only following 9 states on two nearest sites in the wave function (21)

$$\begin{aligned} &|22\rangle, \quad |00\rangle, \quad |20 + 02\rangle, \quad |\uparrow\downarrow - \downarrow\uparrow\rangle, \quad |20 - 02\rangle + c^2 |\uparrow\downarrow + \downarrow\uparrow\rangle, \\ &|\uparrow 0 + 0 \uparrow\rangle, \quad |\uparrow 2 + 2 \uparrow\rangle, \quad |\downarrow 0 + 0 \uparrow\rangle, \quad |\downarrow 2 + 2 \downarrow\rangle \end{aligned} \quad (23)$$

The Hamiltonian for this model has the form which is similar to the previous one given by Eq.(13):

$$\begin{aligned}
H &= \sum_{i=1}^N h_{i,i+1} \\
h_{1,2} &= 1 - 4\eta_1 \cdot \eta_2 + 4\left(1 - \frac{3}{c^4}\right) S_1^z S_2^z + \frac{4}{c^4} \mathbf{S}_1 \cdot \mathbf{S}_2 \\
&- \sum_{\sigma} (c_{1,\sigma}^+ c_{2,\sigma} + c_{2,\sigma}^+ c_{1,\sigma}) (1 - n_{1,-\sigma} - n_{2,-\sigma}) \\
&+ \frac{2}{c^2} \sum_{\sigma} \sigma (c_{1,\sigma}^+ c_{2,\sigma} + c_{2,\sigma}^+ c_{1,\sigma}) (n_{1,-\sigma} - n_{2,-\sigma})^2
\end{aligned} \tag{24}$$

This Hamiltonian for $c > 1$ is also a non-negatively defined operator and Ψ_0 is the exact ground state wave function with zero energy. This Hamiltonian commutes with η^2 , but does not commute with S^2 . Therefore, the eigenfunctions of the Hamiltonian (24) can be described by quantum numbers η and η^z . Making up the same analysis as for previous model we find that for the cyclic model (24) the states with three different values of η have zero energy (as it was for model (13)). They are: one state with $\eta = 0$ and momentum $p = \pi$ (21), all states with $\eta = N/2$ and $p = 0$:

$$\Psi_{N/2, \eta^z} = (\eta^+)^{N/2-\eta^z} |0\rangle \tag{25}$$

and the states with $\eta = N/2 - 1$ and $p = 0$:

$$\Psi_{N/2-1, \eta^z} = (\eta^+)^{N/2-1-\eta^z} \sum_{i < j} (c_{i,\uparrow}^+ c_{j,\downarrow}^+ - c_{i,\downarrow}^+ c_{j,\uparrow}^+) |0\rangle \tag{26}$$

Therefore, for the case of one electron per site ($\eta^z = 0$) the ground state of the model (24) is three-fold degenerate.

The correlation functions in the ground states (25) and (26) obviously coincide with each other in the thermodynamic limit and for half filling case ($\eta^z = 0$) they are:

$$\begin{aligned}
\langle c_{i,\sigma}^+ c_{i+l,\sigma} \rangle &= O\left(\frac{1}{N}\right) & \langle \mathbf{S}_i \mathbf{S}_{i+l} \rangle &= O\left(\frac{1}{N^2}\right) \\
\langle \eta_i^z \eta_{i+l}^z \rangle &= O\left(\frac{1}{N}\right) & \langle \eta_i^- \eta_{i+l}^+ \rangle &= \frac{1}{4} + O\left(\frac{1}{N}\right)
\end{aligned} \tag{27}$$

The existence of ODLRO immediately follows from the form of the wave functions (25) and (26).

The correlation functions in the ground state (21) have similar forms as in Eqs.(16)–(19), and in the thermodynamic limit they reduce to

$$\begin{aligned}
\langle c_{i,\sigma}^+ c_{i+l,\sigma} \rangle &= O\left(\frac{1}{N}\right), & \langle \mathbf{S}_i \mathbf{S}_{i+l} \rangle &= O\left(\frac{1}{N^2}\right), \\
\langle \eta_i^- \eta_{i+l}^+ \rangle &= 2 \langle \eta_i^z \eta_{i+l}^z \rangle = \frac{1}{6} \cos\left(\frac{2\pi l}{N}\right)
\end{aligned} \tag{28}$$

The giant spiral ordering in the last equation implies the existence of ODLRO and, therefore, the superconductivity [14] in the ground state (21). We note that though all




Figure 1: The two-leg ladder

three ground states of the model (24) are superconducting, the properties of these wave functions are essentially different. Let us consider density – density correlator $\langle n_i n_{i+l} \rangle$. For the wave functions (25) and (26) in the thermodynamic limit this correlator decouples: $\langle n_i n_{i+l} \rangle = \langle n_i \rangle \langle n_{i+l} \rangle = 1$. But for the wave function (21) it equals to $\langle n_i n_{i+l} \rangle = 1 + \frac{1}{3} \cos\left(\frac{2\pi l}{N}\right)$.

It is interesting to note that another model having the ground state wave function (21) with $c = 0$ and the same spiral ODLRO (28) can be obtained from the model (1) by simple replacing of operators S by η :

$$H = - \sum_{i=1}^N \eta_i \cdot \eta_{i+1} + \frac{1}{4} \sum_{i=1}^N \eta_i \cdot \eta_{i+2} \quad (29)$$

The direct analogy of this model to the spin model (1) results in the conclusion that the model (29) describes the boundary point on the phase diagram between superconducting and non – superconducting phases where the off-diagonal long-range order is destroyed. We suppose that the model (24) also describes such a point. Thus, the wave functions (9) and (21) are the ground states for the 1D electronic systems in the boundary points between the phases with and without long-range order (ferromagnetic for (13) and off-diagonal for (24)). And we suggest the formation of the ground state with the long-range spiral order like (20) and (28) as a probable scenario of the subsequent destruction of the ferromagnetism and the superconductivity.

The proposed form of wave function can be further generalized for the electronic ladder model (Fig.1). The wave function for the cyclic ladder model containing $2N$ sites has the form:

$$\Psi_0 = P_{S=\eta=0} \langle 0 | g_1 \otimes g_2 \otimes \dots \otimes g_N | 0 \rangle \quad (30)$$

where each g_i corresponds to i -th rung of the ladder:

$$\begin{aligned} g_i = & c_1 \left(a^+ (2x - a^+ a) |\uparrow\uparrow\rangle_i - a |\downarrow\downarrow\rangle_i + (a^+ a - x) |\uparrow\downarrow + \downarrow\uparrow\rangle_i \right) + c_2 |\uparrow\downarrow - \downarrow\uparrow\rangle_i \\ & + c_3 \left(b^+ (2y - b^+ b) |22\rangle_i - b |00\rangle_i + (b^+ b - y) |20 + 02\rangle_i \right) + c_4 |20 - 02\rangle_i \end{aligned} \quad (31)$$

with a^+ , a and b^+ , b are the Bose operators, $|0\rangle$ in (30) is the Bose vacuum of a^+ and b^+ particles, and c_i , x , y are the parameters of the model. $P_{\eta=S=0}$ is the projector onto the state with $S = \eta = 0$.

Let us first consider the case $c_3 = c_4 = 0$. In this case the wave function (30) describes the spin ladder model depending on the two parameters c_2/c_1 and x . It can be shown

that this model coincides with that considered in [12]. It has the singlet ground state (30) degenerate with ferromagnetic state. The spin correlators in the singlet ground state show double-spiral ordering with small shift angle $\Delta\varphi = \frac{2\pi}{N} \frac{2c_2}{c_1}$ between the two giant spirals formed on two legs of the ladder:

$$\begin{aligned}\langle \mathbf{S}_n \mathbf{S}_{n+2l} \rangle &= \frac{1}{4} \cos \left(\frac{2\pi l}{N} \right) \\ \langle \mathbf{S}_n \mathbf{S}_{n+2l+1} \rangle &= \frac{1}{4} \cos \left(\frac{2\pi l}{N} + (-1)^n \Delta\varphi \right)\end{aligned}\quad (32)$$

For the cases of integer or half-integer $x = j$, which correspond to the special cases of the model [12], one can easily recognize in Eq.(31) the Maleev's boson representation of spin $S = j$ operators:

$$S^+ = a^+(2j - a^+a), \quad S^- = a, \quad S^z = a^+a - j$$

Therefore, in these special cases the infinite matrices formed by the Bose operators a^+ , a can be broken off to the size $n = 2j + 1$ and the wave function (30) is reduced to the usual MP form. The spin correlators in the special cases have the exponential decay.

Now let us come back to the general case of the electronic ladder model (30). In order to find the Hamiltonian for which (30) is the exact ground state wave function, one should consider what states are present in the Ψ_0 on the two nearest rungs of the ladder. There are only 26 states from the total 256 states in the product $g_i \otimes g_{i+1}$. Therefore, the Hamiltonian of the ladder model can be written as the sum of the projectors onto the 230 missing states $|\varphi_k\rangle$ with arbitrary positive coefficients λ_k :

$$H = \sum_{i=1}^N h_{i,i+1}, \quad h_{i,i+1} = \sum_{k=1}^{230} \lambda_k |\varphi_k\rangle \langle \varphi_k| \quad (33)$$

Unfortunately, we can not give the explicit form like (13) and (24) for this Hamiltonian, because it has a very cumbersome form. But we are able to determine some properties of the Hamiltonian (33). This Hamiltonian commutes with both S^2 and η^2 . It has multiply degenerated ground state: the state with $S = \eta = 0$ (30) and all the states with $S + \eta = N$ have zero energy. Hence the electronic ladder model (33) describes the boundary point between the phases with and without ferromagnetic and off-diagonal long-range order. The correlation functions in the ground state (30) can be calculated with the use of the technique developed in Appendix. For the case $c_1 > c_3$ there is the same double-spiral spin ordering (32) as for the spin ladder model, while all other correlations are exponentially small. For the case $c_1 < c_3$ the double-spiral ODLRO is realized. In the most interesting symmetric case $c_1 = c_3$, $c_2 = c_4$, $x = y$ the system possesses both the giant spiral spin order and the giant spiral ODLRO:

$$\begin{aligned}\langle \mathbf{S}_n \mathbf{S}_{n+2l} \rangle &= \langle \eta_n \eta_{n+2l} \rangle = \frac{1}{8} \cos \left(\frac{2\pi l}{N} \right) \\ \langle \mathbf{S}_n \mathbf{S}_{n+2l+1} \rangle &= \langle \eta_n \eta_{n+2l+1} \rangle = \frac{1}{8} \cos \left(\frac{2\pi l}{N} + (-1)^n \Delta\varphi \right)\end{aligned}\quad (34)$$

where $\Delta\varphi = \frac{2\pi}{N} \frac{2c_2}{c_1}$.

Therefore, in this case the wave function (30) describes the boundary points on the phase diagram between the four different phases: the singlet phases with and without ODLRO, and the ferromagnetic phases with and without ODLRO. In the special cases when x (y) is integer or half-integer the spin (off-diagonal) correlations exponentially decay and the wave function in the corresponding Bose space can be represented in the MP form with finite matrices of the size $n = 2x + 1$ or $n = 2y + 1$. When both x and y are integers or half-integers the wave function (30) can be written in the usual MP form with the size of matrices $n = (2x + 1)(2y + 1)$.

4 Summary

We have found another form of the singlet ground state wave function for the quantum spin model considered previously in [9, 10]. The special technique was developed for the exact calculation of the norm and the correlation functions. This form of the wave function allowed us to generalize it for two 1D electronic models.

The first model describes the half-filling electronic system in the ferromagnet – antiferromagnet transition point when the singlet and the ferromagnetic states are degenerate. In the singlet ground state the spin correlators show the giant spiral magnetic ordering with the period of the spiral equals to the system size, while all other correlations vanish in the thermodynamic limit.

The second electronic model in the half filling case has three-fold degenerate ground state. All ground states have off-diagonal long-range order and, therefore, are superconducting. The calculation of the correlation functions shows that one of the ground states has a giant spiral ODLRO.

The comparison of these electronic models with original spin model [11, 12] leads us to the conclusion that these two electronic models describe the boundary points on the phase diagram between the phases with and without long-range order (ferromagnetic for the first and off-diagonal for the second model). Therefore, we presume that if the Hamiltonian of the 1D quantum system commutes with operators forming the $SU(2)$ algebra (it can be the spin S or the pseudo-spin η operators), then the appearance of the ground state with the giant spiral order predicts the following destruction of the ferromagnetism or the superconductivity.

We have briefly considered the generalization of the proposed form of the wave function for the electronic ladder model. The general case of this model has much more rich phase diagram than two first models. In some particular cases this model describes the boundary points on the phase diagram between four different phases: with and without ferromagnetic and off-diagonal long-range order. There are also some special cases of the electronic ladder model when the ground state wave function is reduced to the usual MP form. Besides, the proposed form of the wave function can be also generalized for the 2D case and different types of lattices.

5 Acknowledgements

Authors are grateful to Max-Planck-Institut für Physik Komplexer Systeme for kind hospitality. This work was supported by RFFR.

Appendix

Let us calculate the norm of the wave function Ψ_0 (2)

$$\langle \Psi_0 | \Psi_0 \rangle = \langle \Psi | P_0 | \Psi \rangle \quad (\text{A.1})$$

Since the function Ψ has $S^z = 0$, then the projector P_0 in the Eq.(4) takes the form [10]

$$P_0 = \frac{1}{2} \int_0^\pi \sin \gamma d\gamma e^{izS^-} e^{iz'S^+} \quad (\text{A.2})$$

where $z = \tan \frac{\gamma}{2}$, $z' = \sin \frac{\gamma}{2} \cos \frac{\gamma}{2}$ and $S^{+(-)}$ are the operators of the total spin.

Therefore, the norm takes the form:

$$\langle \Psi_0 | \Psi_0 \rangle = \frac{1}{2} \int_0^\pi \sin \gamma d\gamma \langle 0_a, 0_b | \prod_{i=1}^N (g_i^+ e^{izS_i^-} e^{iz'S_i^+} g_i) | 0_a, 0_b \rangle$$

where

$$\begin{aligned} g_i^+ e^{izS_i^-} e^{iz'S_i^+} g_i &= (a^+ |\uparrow_i| + a |\downarrow_i|) e^{izS_i^-} e^{iz'S_i^+} (b^+ |\uparrow_i| + b |\downarrow_i|) \\ &= a^+ b^+ + (1 - z'z)ab + izab^+ + iz'a^+b \end{aligned}$$

and a^+ , a are the Bose operators.

So, the norm can be rewritten as follows

$$\langle \Psi_0 | \Psi_0 \rangle = \frac{1}{2} \int_0^\pi \sin \gamma d\gamma \langle 0 | G^N | 0 \rangle \quad (\text{A.3})$$

with $|0\rangle = |0_a, 0_b\rangle$ is the Bose vacuum of a^+ and b^+ particles and

$$G = u(a^+ b^+ + ab) + iv(ab^+ + a^+ b)$$

where $u = \cos \frac{\gamma}{2}$, $v = \sin \frac{\gamma}{2}$.

Let us introduce the auxiliary function $P(\xi)$:

$$P(\xi) = \langle 0 | e^{\xi G} | 0 \rangle \quad (\text{A.4})$$

then,

$$\langle 0 | G^N | 0 \rangle = \left. \frac{d^N P}{d\xi^N} \right|_{\xi=0}$$

In order to find $P(\xi)$ we perform the following manipulations. First, we take the derivative of $P(\xi)$:

$$\frac{dP}{d\xi} = \langle 0 | G e^{\xi G} | 0 \rangle = \langle 0 | e^{\xi G} G | 0 \rangle = u \langle 0 | ab e^{\xi G} | 0 \rangle = u \langle 0 | e^{\xi G} a^+ b^+ | 0 \rangle \quad (\text{A.5})$$

Now we need to carry $a^+ b^+$ over G in the last expression

$$e^{\xi G} a^+ b^+ = e^{\xi G} a^+ e^{-\xi G} e^{\xi G} b^+ e^{-\xi G} e^{\xi G} \quad (\text{A.6})$$

It can be made by using the following equations:

$$\begin{aligned}
e^{\xi G} a^+ e^{-\xi G} &= a^+ \cos \xi + (ub + ivb^+) \sin \xi \\
e^{\xi G} a e^{-\xi G} &= a \cos \xi - (ub^+ + ivb) \sin \xi \\
e^{\xi G} b^+ e^{-\xi G} &= b^+ \cos \xi + (ua + iva^+) \sin \xi \\
e^{\xi G} b e^{-\xi G} &= b \cos \xi - (ua^+ + iva) \sin \xi
\end{aligned} \tag{A.7}$$

Substituting Eqs.(A.6), (A.7) into Eq.(A.5), we find

$$\langle 0|e^{\xi G} a^+ b^+|0\rangle = u \sin \xi \cos \xi \langle 0|e^{\xi G}|0\rangle + u^2 \sin^2 \xi \langle 0|ab e^{\xi G}|0\rangle$$

The last equation can be rewritten as the differential equation on the $P(\xi)$

$$\frac{dP}{d\xi} = u^2 \sin \xi \cos \xi P(\xi) + u^2 \sin^2 \xi \frac{dP}{d\xi} \tag{A.8}$$

with boundary condition $P(0) = 1$.

The solution of Eq.(A.8) is

$$P(\xi) = \frac{1}{\sqrt{1 - u^2 \sin^2 \xi}} \tag{A.9}$$

Integrating Eq.(A.3) over γ , we obtain

$$\langle \Psi_0|\Psi_0\rangle = \frac{1}{2} \int_0^\pi \sin \gamma d\gamma \left. \frac{d^N P}{d\xi^N} \right|_{\xi=0} = \frac{d^N}{d\xi^N} \left(\frac{1}{\cos^2(\frac{\xi}{2})} \right) \Big|_{\xi=0} \tag{A.10}$$

So, finally we arrive at

$$\langle \Psi_0|\Psi_0\rangle = 2 \frac{d^{N+1}}{d\xi^{N+1}} \left(\tan \frac{\xi}{2} \right) \Big|_{\xi=0} = \frac{4(2^{N+2} - 1)}{N + 2} |B_{N+2}| \tag{A.11}$$

Here B_N are the Bernoulli numbers.

To calculate the spin correlators we need to introduce operators:

$$\begin{aligned}
G_z &= g_i^+ e^{izS_i^-} e^{iz'S_i^+} 2S_i^z g_i = u(a^+ b^+ - ab) + iv(ab^+ - a^+ b) \\
G_+ &= g_i^+ e^{izS_i^-} e^{iz'S_i^+} S_i^+ g_i = ua^+ b + ivab \\
G_- &= g_i^+ e^{izS_i^-} e^{iz'S_i^+} S_i^- g_i = uab^+ + iva^+ b^+
\end{aligned}$$

Then, the correlator $\langle \mathbf{S}_1 \mathbf{S}_{l+1} \rangle$ will be defined by

$$\langle \Psi_0|\mathbf{S}_1 \mathbf{S}_{l+1}|\Psi_0\rangle = \frac{1}{2} \int_0^\pi \sin \gamma d\gamma \langle 0| \frac{1}{4} G_z G^l G_z G^{N-l-2} + \frac{1}{2} G_+ G^l G_- G^{N-l-2} |0\rangle \tag{A.12}$$

(since $\langle 0|G_- \dots |0\rangle = 0$).

The expectation values in Eq.(A.12) can be represented as

$$\begin{aligned}
\langle 0| G_z G^l G_z G^{N-l-2} |0\rangle &= \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \langle 0| G_z e^{\xi G} G_z e^{\zeta G} |0\rangle \Big|_{\xi=\zeta=0} \\
\langle 0| G_+ G^l G_- G^{N-l-2} |0\rangle &= \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \langle 0| G_+ e^{\xi G} G_- e^{\zeta G} |0\rangle \Big|_{\xi=\zeta=0}
\end{aligned} \tag{A.13}$$

After the procedure similar to that for the norm and the integration over γ , we obtain

$$\langle \Psi_0|\mathbf{S}_1 \mathbf{S}_{l+1}|\Psi_0\rangle = \frac{\partial^l}{\partial \xi^l} \frac{\partial^{N-l-2}}{\partial \zeta^{N-l-2}} \left(-\frac{3 \cos(\xi - \zeta)}{8 \cos^4(\frac{\xi+\zeta}{2})} \right) \Big|_{\xi=\zeta=0} \tag{A.14}$$

References

- [1] M.Fannes, B.Nachtergaele, R.F.Werner, Commun. Math. Phys. **144**, 443 (1992).
- [2] A. Klumper, A. Schadschneider, J. Zittartz, Z. Phys. B. **87**, 281 (1992); Europhys. Lett. **24**(4), 293 (1993).
- [3] A.K.Kolezhuk, H.-J.Mikeska, Phys. Rev. Lett. **80**, 2709 (1998); Int. J. Mod. Phys. **B12**, 2325 (1998).
- [4] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, Phys. Rev. Lett. **59**, 799 (1987); Commun. Math. Phys. **115**, 477 (1988).
- [5] D.V.Dmitriev, V.Ya.Krivnov, A.A.Ovchinnikov, JETP **88**, 138 (1999).
- [6] R.Strack, D.Vollhardt, Phys. Rev. Lett. **70**, 2637 (1993), Phys. Rev. Lett.**72**, 3425 (1994).
- [7] A.A.Ovchinnikov, Mod. Phys. Lett B**7**, 1397 (1993); J. Phys. **CM 6**, 11057 (1994).
- [8] Jan de Boer, A.Schadschneider, Phys. Rev. Lett. **75**, 4298 (1995).
- [9] T.Hamada, J.Kane, S.Nakagawa and Y.Natsume. J. Phys. Soc. Jpn. **57**, 1891 (1988); **58**, 3869 (1989).
- [10] D.V.Dmitriev, V.Ya.Krivnov, A.A.Ovchinnikov, Z. Phys. B**103** 193, (1997); Phys. Rev. B**56**, 5985 (1997).
- [11] V.Ya.Krivnov, A.A.Ovchinnikov, Phys. Rev. B**53**, 6435 (1996).
- [12] D.V.Dmitriev, V.Ya.Krivnov, A.A.Ovchinnikov (to be published).
- [13] P.van Leuven, Physica **45**, 86 (1969).
- [14] C.N.Yang, Rev. Mod. Phys. **34**, 694 (1962).
- [15] C.N.Yang, Phys. Rev. Lett. **63**, 2144 (1989).
- [16] F.H.L.Essler, V.E.Korepin, K.Schoutens, Phys. Rev. Lett. **68**, 2960 (1992); Phys. Rev. Lett. **70**, 73 (1993).